

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) April 2012		2. REPORT TYPE Viewgraph		3. DATES COVERED (From - To) April 2012- June 2012	
4. TITLE AND SUBTITLE Volatilization mechanism of 1-ethyl-3-methylimidazolium bromide ionic liquid			5a. CONTRACT NUMBER In-House		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)  G.L. Vaghjiani			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER 23031166		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/RQRP 10 E.Saturn Blvd. Edwards AFB CA 93524-7680			8. PERFORMING ORGANIZATION REPORT NO.		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB CA 93524-7048			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RZ-ED-VG-2012-200		
12. DISTRIBUTION / AVAILABILITY STATEMENT Distribution A: Approved for Public Release; Distribution Unlimited. PA#12466					
13. SUPPLEMENTARY NOTES Conference paper for the 22nd International Symposium on Gas Kinetics, Boulder, CO in 18-22 June 2012.					
14. ABSTRACT Although molten salts, more recently known as ionic liquids (ILs), have been known for almost a century (for instance, ethylammonium nitrate, mp =12 °C, was discovered in 1914), the rapid development of ionic liquid chemistry has only occurred within the last decade. In 2000, only 217 papers were published on the subject, whereas in 2010, over 6,000 papers appeared in the literature. The number of possible ILs has been estimated at 10 <sup>18</sup> . Detailed studies of ILs offer insight into new and more complex fundamental chemistry, as well as allow for an assessment for their potential application in emerging technologies such as in solar cells, fuel electrolytes, bio- or nano-catalysts, high-energy-density propellants, and green solvents. In order to better understand the volatilization process for ILs, the vapor evolved from heating the ionic liquid 1-ethyl-3-methylimidazolium bromide (EMIM+Br-) was analyzed via tunable vacuum ultraviolet photoionization time of flight mass spectrometry (VUV-PI-TOFMS) and thermal gravimetric analysis mass spectrometry (TGA-MS). For this ionic liquid, the experimental results indicate that vaporization takes place via the evolution of alkyl bromides and alkylimidazoles, presumably through alkyl abstraction via an SN2 type mechanism, and that vaporization of intact ion pairs or the formation of carbenes is negligible. Activation enthalpies for the formation of the methyl and ethyl bromides were evaluated experimentally, $\square H_{\ddagger}(\text{CH}_3\text{Br}) = 116.1 \pm 6.6$ kJ/mol and $\square H_{\ddagger}(\text{CH}_3\text{CH}_2\text{Br}) = 122.9 \pm 7.2$ kJ/mol, and the results are found to be in agreement with calculated values for the SN2 reactions. Comparisons of product photoionization efficiency (PIE) curves with literature data are in good agreement, and <i>ab initio</i> thermodynamics calculations are presented as further evidence for the proposed thermal decomposition mechanism. Estimates for the enthalpy of vaporization of 1-ethyl-3-methylimidazolium bromide and, by comparison, 1-butyl-3-methylimidazolium bromide (BMIM+Br-) from molecular dynamics calculations and their gas phase enthalpies of formation obtained by G4 calculations yield estimates for the ionic liquids, enthalpies of formation in the liquid phase: $\square H_{\text{vap}}(298\text{ K})(\text{EMIM}+\text{Br}^-) = 168 \pm 20$ kJ/mol, $\square H_{\text{f, gas}}(298\text{ K})(\text{EMIM}+\text{Br}^-) = 38.4 \pm 10$ kJ/mol, $\square H_{\text{f, liq}}(298\text{ K})(\text{EMIM}+\text{Br}^-) = -130 \pm 22$ kJ/mol, $\square H_{\text{f, gas}}(298\text{ K})(\text{BMIM}+\text{Br}^-) = -5.6 \pm 10$ kJ/mol, and $\square H_{\text{f, liq}}(298\text{ K})(\text{BMIM}+\text{Br}^-) = -180 \pm 20$ kJ/mol.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  SAR	18. NUMBER OF PAGES  16	19a. NAME OF RESPONSIBLE PERSON Tommy Hawkins
a. REPORT  Unclassified	b. ABSTRACT  Unclassified	c. THIS PAGE  Unclassified			19b. TELEPHONE NO (include area code) 661-275-5449

# Volatilization Mechanism of 1-Ethyl-3-methylimidazolium Bromide Ionic Liquid

---



**Ghanshyam L. Vaghjiani**

**AFRL/RZSP**

**Propellants Branch**

**Space and Missile Propulsion Division**

**Propulsion Directorate**

**Air Force Research Laboratory**

**AFRL/RZSP**

**10 E Saturn Blvd**

**Edwards AFB, CA 93524**

**Tel: 661 275 5657**

**Fax: 661 275 5471**

**Email: [ghanshyam.vaghjiani@edwards.af.mil](mailto:ghanshyam.vaghjiani@edwards.af.mil)**



# Acknowledgements



AFRL Edwards:  
**Steve Chambreau**  
**Jerry Boatz**



ALS/UC Berkeley:  
**Christine Koh**  
**Oleg Kostko**  
**Amir Golan**  
**Stephen Leone**



AFOSR:  
**Mike Berman**





# Motivation to Study Room Temperature Ionic Liquids




## ☐ Molten Salts @ RT

- ☐ Low mp
- ☐ Large (organic) ions
- ☐ Low volatility ( $vp_{298\text{ K}} \ll 1\text{ Pa}$ )
- ☐ Thermally stable
- ☐ Good conductivity
- ☐ Good solvents

☐ Unique reactivity

☐ Popularized in green/sustainable chemistry applications



- ☐  $> 10^{18}$  cation/anion pair combinations
- ☐ Enormous tunability



# Hypergolic Ignition



- ❑ Autoignition at ambient P & T

- ❑ MMH/NTO current hypergol propellants

- ❑ IL hypergol fuels low vapor toxicity

- ❑ Operational cost savings (cf MMH)

- ❑ Higher performing than MMH

- ❑ Synthesis tunability

- ❑  $T_b$  (calculated)  $\gg T_d$  for ILs

- ❑ What role does decomposition/vaporization play in understanding IL/ox autoignition?




JPC-A 2008, 112, 7816






# How Do ILs Volatilize?





- ☐ Understand gas phase structure of ILs
- ☐ Determine heat of vaporization..... calibrate MD simulations
- ☐ (for  $\Delta H_{f(l)}$  and therefore  $I_{sp}$ )
- ☐ Characterize thermal decomposition of ILs  
(basicity & nucleophilicity of anion)





## Separate temperature control of reservoir and outlet





7.4-15.0 eV photons, 0.025 eV resolution,  
0.040 eV absolute accuracy



# Structure of Aprotic ILs in the Gas Phase



- PIE curves and structures



- Dissociative photoionization



JPC-A 2010, 114, 879


Appearance energies are:

[emim][Tf<sub>2</sub>N]: 8.72 ± 0.03 eV  
[emim][Pf<sub>2</sub>N]: 8.78 ± 0.04 eV  
[dmpim][Tf<sub>2</sub>N]: 8.59 ± 0.03 eV





# Volatilization of EMIM<sup>+</sup>Br<sup>-</sup>



☐ No ion signal at 111


☐ No gas phase IL ion pairs

☐ No dissociative ionization of IL ion pairs

☒ New process for volatilization



# Photoionization Energetics of Gas Phase EMIM<sup>+</sup>Br<sup>-</sup>



Is this responsible for 94 & 96 ion masses?

Is this responsible for 108 & 110 ion masses?

Is this responsible for 82 ion mass?

Is this responsible for 96 ion mass?



No 111 ion mass

How would you go about detecting this channel?

MP2/6-31+G(d,p), 0 K, ZPVE corrected



# Photoionization Efficiency (PIE) Curves





Ref 60:

Baer, T.; Song, Y.; Liu, J.; Chen, W.; Ng, C. Y. *Faraday Discuss.* **2000**, 115, 137


Ref 61:

Locht, R.; Leyh, B.; Dahareng, D.; Hottmann, K.; Jochims, H. W.; Baumgartel, H. *Chem. Phys.* **2006**, 323, 458





# Photoionization Efficiency (PIE) Curves




All experimental AEs consistent with MP2 values for dissociative photoionization of IL ion pair but no 111 ion mass

Is there another explanation?





# Thermolysis of EMIM+Br<sup>-</sup> can Explain observed PIE curves






Observed AEs consistent with experimental/MP2/6-31+G(d,p) IEs

VUV beamline is sampling thermolysis not vaporization products





# TGA/MS Analysis of EMIM<sup>+</sup>Br<sup>-</sup> Thermolysis







# $\Delta H_{\text{vap}}$ for EMIM<sup>+</sup>Br<sup>-</sup> & BMIM<sup>+</sup>Br<sup>-</sup>



MD values


	T (K)	$\Delta H_{\text{vap}}$ (kJ/mol)
EMIM <sup>+</sup> Br <sup>-</sup>	298	147.1 <sup>b</sup>
EMIM <sup>+</sup> Br <sup>-</sup>	393	138.3 <sup>a</sup>
EMIM <sup>+</sup> Br <sup>-</sup>	473	130.8 <sup>a</sup>
BMIM <sup>+</sup> Br <sup>-</sup>	298	152.8 <sup>b</sup>
BMIM <sup>+</sup> Br <sup>-</sup>	323	150.5 <sup>a</sup>
BMIM <sup>+</sup> Br <sup>-</sup>	393	144.0 <sup>a</sup>

<sup>a</sup> MD direct calculation. <sup>b</sup> Linear extrapolation from higher temperature MD values.

298 K values, this study

$$\begin{aligned}\Delta H_{\text{vap}}(\text{EMIM}^+\text{Br}^-) &= 168 \pm 20 \text{ kJ/mol} \\ \Delta H_{\text{f, gas}}(\text{EMIM}^+\text{Br}^-) &= 38.4 \pm 10 \text{ kJ/mol} \\ \Delta H_{\text{f, liq}}(\text{EMIM}^+\text{Br}^-) &= -130 \pm 22 \text{ kJ/mol} \\ \Delta H_{\text{f, gas}}(\text{BMIM}^+\text{Br}^-) &= -5.6 \pm 10 \text{ kJ/mol} \\ \Delta H_{\text{f, liq}}(\text{BMIM}^+\text{Br}^-) &= -180 \pm 20 \text{ kJ/mol}\end{aligned}$$

Calorimetric / *ab initio* values



Paulechka, Y. U.; Kabo, A. G.; Blokhin, A. V. *J. Phys. Chem. B* **2009**, 113, 14742



# Conclusions



- ☐ **VUV PI ToF MS spectroscopy has provided**
  - ☐ **Structural information on volatilized species from ILs**
  - ☐ **AEs & PIE curves, which leads to unique identification**
  - ☐  **$\Delta H_{\text{vap}}$ , which leads to  $\Delta H_{\text{liq}}$  & allows MD calibrations**
  - ☐ **Insight into the dynamics of dissociative photoionization**
  
- ☐ **Nature of the IL volatiles are cation/anion dependent**
  - ☐ **Ion pairs**
  - ☐ **Neutrals from proton transfer**
  - ☐ **Neutrals from decomposition**
  
- ☐ **IL combustion is influenced by the nature of the volatiles (fuel)**